

# Publicly Funded Immunization Schedules for Ontario – January 2021

Publicly funded vaccines may be provided only to eligible individuals and must be free of charge

## Routine Schedule: Children Starting Immunization in Infancy

Vaccine	Age	2 Months	4 Months	6 Months	1 Year <sup>Φ</sup>	15 Months	18 Months	4 Years	Grade 7	14 Years	24 Years	≥34 Years <sup>Υ</sup>	65 Years	
<b>DTaP-IPV-Hib</b> Diphtheria, Tetanus, Pertussis, Polio, <i>Haemophilus influenzae</i> type b		◆	◆	◆			◆							
<b>Pneu-C-13</b> Pneumococcal Conjugate 13		◆	◆		◆									
<b>Rot-5</b> Rotavirus		▲	▲	▲										
<b>Men-C-C</b> Meningococcal Conjugate C					◆									
<b>MMR</b> Measles, Mumps, Rubella					■									
<b>Var</b> Varicella						■								
<b>MMRV</b> Measles, Mumps, Rubella, Varicella								■						
<b>Tdap-IPV</b> Tetanus, diphtheria, pertussis, Polio								◆						
<b>HB</b> Hepatitis B									●					
<b>Men-C-ACYW</b> Meningococcal Conjugate ACYW-135									●					
<b>HPV-9</b> Human Papillomavirus									●					
<b>Tdap</b> Tetanus, diphtheria, pertussis										◆	◆			
<b>Td (booster)</b> Tetanus, diphtheria												◆ Every 10 years		
<b>HZ</b> Herpes Zoster													I	
<b>Pneu-P-23</b> Pneumococcal Polysaccharide 23													■	
<b>Inf</b> Influenza													*Every year in the fall	

- ◆ - A single vaccine dose given by intramuscular injection
- - A single vaccine dose given by subcutaneous injection
- ▲ - A single vaccine dose given by mouth
- - Provided through school-based immunization programs. Men-C-ACYW is a single dose; HB is a 2 dose series (see Table 6); HPV-9 is a 2 dose series (see Table 10). Each vaccine dose is given by intramuscular injection
- Φ - Given no earlier than the 1<sup>st</sup> birthday, and prior to 16 months of age

- Υ - Once a dose of Tdap is given in adulthood (24 years of age), adults should receive Td boosters every 10 years thereafter
  - I - HZ is a 2 dose series (see Table 12) given by intramuscular injection
  - \* - Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given ≥4 weeks apart. Children who have previously received ≥1 dose of influenza vaccine should receive 1 dose per season thereafter
- Note:** A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)

# Publicly Funded Immunization Schedules for Ontario – January 2021

Publicly funded vaccines may be provided only to eligible individuals and must be free of charge

Catch-up Schedule 1: Children Starting Immunization between 1-6 Years																				
Vaccine	Age	1 <sup>st</sup> Visit:			2 <sup>nd</sup> Visit: 2 months after 1 <sup>st</sup> visit					3 <sup>rd</sup> Visit: 2 months after 2 <sup>nd</sup> visit		4 <sup>th</sup> Visit: 6-12 months after 3 <sup>rd</sup> visit		5 <sup>th</sup> Visit (only required if child was <4 years at 4 <sup>th</sup> visit): 4-6 yrs of age and 6-12 months after 4 <sup>th</sup> visit	Grade 7	14-18 yrs <sup>§</sup>	24-28 yrs		≥34 yrs <sup>¶</sup>	65 yrs
		If child is			If child is <5 years and was			If child is		If child is		If adult was								
		<4 yrs	4 yrs	5-6 yrs	<2 yrs at 1 <sup>st</sup> visit	2-3 yrs at 1 <sup>st</sup> visit	4 yrs at 1 <sup>st</sup> visit	5-6 yrs	7 yrs	<7 yrs	7 yrs	<4 yrs	4-8 yrs				<18 yrs at previous visit <sup>†</sup>	≥18 yrs at previous visit <sup>¶</sup>		
DTaP-IPV-Hib		◆	◆	◆	◆	◆	◆			◆		◆								
Pneu-C-13		◆	◆		◆															
MMR		■																		
MMRV			■	■									■	■						
Var					■	■														
Men-C-C		◆	◆	◆																
Tdap-IPV									◆	◆		◆	◆							
HB															●					
Men-C-ACYW															●					
HPV-9															●					
Tdap																◆	◆			
Td																		◆	◆	◆ Every 10 years
HZ																				┆
Pneu-P-23																				■
Inf		*Every year in the fall																		

◆ - A single vaccine dose given by intramuscular injection

■ - A single vaccine dose given by subcutaneous injection

● - Provided through school-based immunization programs. Men-C-ACYW is a single dose; HB is a 2 dose series (see Table 6); HPV-9 is a 2 dose series (see Table 10). Each vaccine dose is given by intramuscular injection

§ - Given 10 years after the (4-8 year old) Tdap-IPV dose

† - Given 10 years after the adolescent Tdap dose

¶ - Once a dose of Tdap is given in adulthood (>18 yrs), adults should receive Td boosters every 10 years thereafter

┆ - HZ is a 2 dose series (see Table 12) given by intramuscular injection

\* - Children 6 months to 8 years of age who have not previously received a dose of influenza vaccine require 2 doses given ≥4 weeks apart. Children who have previously received ≥1 dose of influenza vaccine should receive 1 dose per season thereafter

**Note:** A different schedule and/or additional doses may be needed for high risk individuals (see Table 3) or if doses of a vaccine series are missed (see appropriate Tables 4-24)